

**UF** | UNIVERSITY of
FLORIDA

Machine-Learning-Based Data Reconstruction Chain for the Short Baseline Near Detector

B. Carlson


SBND

July 1, 2025

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Workforce Development for Teachers and Scientists, Office of Science Graduate Student Research (SCGSR) program. The SCGSR program is administered by the Oak Ridge Institute for Science and Education for the DOE under contract number DE-SC0014664.

Neutrinos



- Neutrinos are electrically neutral, nearly massless leptons with three flavor generations



	mass → $\approx 2.3 \text{ MeV}/c^2$ charge → $2/3$ spin → $1/2$ u up	mass → $\approx 1.275 \text{ GeV}/c^2$ charge → $2/3$ spin → $1/2$ c charm	mass → $\approx 173.07 \text{ GeV}/c^2$ charge → $2/3$ spin → $1/2$ t top	mass → 0 charge → 0 spin → 1 g gluon	mass → $\approx 126 \text{ GeV}/c^2$ charge → 0 spin → 0 H Higgs boson	
QUARKS	mass → $\approx 4.8 \text{ MeV}/c^2$ charge → $-1/3$ spin → $1/2$ d down	mass → $\approx 95 \text{ MeV}/c^2$ charge → $-1/3$ spin → $1/2$ s strange	mass → $\approx 4.18 \text{ GeV}/c^2$ charge → $-1/3$ spin → $1/2$ b bottom	mass → 0 charge → 0 spin → 1 γ photon		
	mass → $0.511 \text{ MeV}/c^2$ charge → -1 spin → $1/2$ e electron	mass → $105.7 \text{ MeV}/c^2$ charge → -1 spin → $1/2$ μ muon	mass → $1.777 \text{ GeV}/c^2$ charge → -1 spin → $1/2$ τ tau	mass → $91.2 \text{ GeV}/c^2$ charge → 0 spin → 1 Z Z boson	GAUGE BOSONS	
	mass → $< 2.2 \text{ eV}/c^2$ charge → 0 spin → $1/2$ ν_e electron neutrino	mass → $< 0.17 \text{ MeV}/c^2$ charge → 0 spin → $1/2$ ν_μ muon neutrino	mass → $< 15.5 \text{ MeV}/c^2$ charge → 0 spin → $1/2$ ν_τ tau neutrino	mass → $80.4 \text{ GeV}/c^2$ charge → ± 1 spin → 1 W W boson		
LEPTONS						

Neutrinos




- Neutrino interact in flavor states (*interaction basis*) via weak force bosons
 - Charged current (CC) interact via W^\pm bosons
 - Neutral current (NC) interact via Z^0 boson



Neutrinos


- Neutrino flavor states (*interaction basis*) are mixed between mass states (*mass basis*)
 - This gives rise to **neutrino oscillations**

$$|\nu_\alpha\rangle = \sum_{i=1}^3 U_{\alpha i}^* |\nu_i\rangle$$



$$P_{\nu_\alpha \rightarrow \nu_\beta}(L) = |\langle \nu_\beta | \nu_\alpha(L) \rangle|^2 = \left| \sum_j U_{\alpha j}^* U_{\beta j} e^{-im_j^2 L / (2E)} \right|^2$$

Neutrinos



What are the neutrino masses?

*Beta decay experiments e.g. **KATRIN***

*Cosmology e.g. **Planck (CMB + BAO)***

Are neutrinos their own antiparticles (Majorana vs. Dirac)?

*Neutrinoless double beta decay e.g. **LEGEND, CUORE, NEXT, nEXO, KamLAND-Zen***

Is there CP violation in the neutrino sector?


*Long baseline experiments e.g. **T2K, NOvA, DUNE, Hyper K etc.***

Are there more than three neutrinos?

*Short baseline experiments e.g. **LSND, MiniBooNE, SBN***


Neutrinos

- 3 active neutrinos from measuring the Z^0 boson branching ratio
- Leaves the possibility of one or more *sterile neutrinos*, which do not interact with the weak nuclear force



LSND ν_e excess at low energy

4.8 σ





MiniBooNE ν_e excess at low energy

Short-Baseline Neutrino (SBN) Program



P. Machado, O. Palamara, D. Schmitz
Ann.Rev.Nucl.Part.Sci. 69 (2019) 363-387

Short-Baseline Neutrino Program at Fermilab (2024)



SBN program and detectors



$$P(\nu_\mu \rightarrow \nu_e) \simeq \sin^2(2\theta_{\mu e}) \sin^2\left(\frac{\Delta m_{41}^2 L}{4E}\right)$$


$$\sin^2(2\theta_{\mu e}) \equiv 4 |U_{e4} U_{\mu 4}|^2$$




Short-Baseline Neutrino (SBN) Program

- Compute **expected number of events** as a function of E_ν
- O(10-15%) uncertainty on **neutrino xsec** and **flux** and E_ν **resolution**

$$N(E_\nu) = B + \int P_{\nu_\mu \rightarrow \nu_e}(E_\nu) \times \Phi(E_\nu) \times \sigma(E_\nu) \times n_t \times \epsilon(E_\nu) \times D(E_{\text{reco}}, E_\nu) dE_\nu$$




MicroBooNE only result




Projected full SBN sensitivity assuming oscillations

Short-Baseline Neutrino (SBN) Program




- Short Baseline Near Detector (SBND) - Constrain flux and cross section systematics
- ICARUS - Measure neutrino oscillations by taking ratio of events at near and far detectors
- Both detectors are **Liquid Argon Time Projection Chambers (LArTPC)**

Liquid Argon Time Projection Chamber (LArTPC)




SBND Technical Design

Liquid Argon Time Projection Chamber (LArTPC)



SBND Technical Design


Liquid Argon Time Projection Chamber (LArTPC)



SBND Technical Design

Liquid Argon Time Projection Chamber (LArTPC)


- Precise tracking
- Cheap ($O(1)$ \$/kg)
- Dense (1.4 g/cm^3)
- Detailed calorimetry
- High A
- Scalable



SBND Technical Design

Short Baseline Near Detector (SBND)

- SBND is a monolithic LArTPC with 2 TPCs divided by a cathode plane assembly (CPA) [1]
- 3 wireplanes with 2 induction wireplanes at ± 60 deg
 - **3 mm** wire and inter-plane spacing
 - 11,892 wires per anode plane assembly (APA)
- 120 photomultiplier tubes (PMTs) and 192 X-ARAPUCAs* [2]




SBND with coordinate system

[1] Construction of precision wire readout planes for the Short-Baseline Near Detector (SBND) [arXiv:2002.08424](https://arxiv.org/abs/2002.08424)

[2] Scintillation Light in SBND: Simulation, Reconstruction, and Expected Performance of the Photon Detection System [arXiv:2406.07514](https://arxiv.org/abs/2406.07514)

*Silicon-based photomultiplier light trap


SBND circa. March 2023





Short Baseline Near Detector (SBND)



First neutrinos recorded in July, 2024 - [press release](#)





1/2 TPC event display



Short Baseline Near Detector (SBND)


$\nu_\mu - {}^{40}\text{Ar}$
2M ν_μ CC events in 1 year

$\nu_e - {}^{40}\text{Ar}$
15k ν_e CC events in 1 year




- Expect millions neutrino interactions for SBND's lifetime (10e20 POT)
- Collected neutrinos from $>2.57\text{e}20$ POT already


SBND Cumulative POT through May 20, 2025




Liquid Argon Time Projection Chamber (LArTPC)




Liquid Argon Time Projection Chamber (LArTPC)




Liquid Argon Time Projection Chamber (LArTPC)



Liquid Argon Time Projection Chamber (LArTPC)





Liquid Argon Time Projection Chamber (LArTPC)




Energy

- For charged current (CC) interactions, E_ν is the sum of the visible components (E_ℓ, E_h) minus the binding energy of the nucleus E_{nuc}


$$E_\nu = E_\ell + E_h - E_{nuc}$$




Low momentum ← → High momentum



$$\Delta x = \int_0^{E_0} \frac{1}{dE/dx} dE$$



$$\theta_0 = \frac{S_2}{p\beta c} z \sqrt{\frac{x}{X_0}} \left[1 + \epsilon \log \left(\frac{x}{X_0} \right) \right]$$




$$E_{\text{shower}} = \underbrace{W_i}_{[\text{MeV}]} \cdot \underbrace{C_{\text{calo}}}_{[\text{MeV}/e]} \cdot \underbrace{C_{\text{adj}}}_{[e^-/\text{ADC}]} \cdot \frac{1}{R} \cdot \sum_Q e^{\frac{t_{\text{drift}}}{\tau}} Q$$

Calorimetric constant (TPC gain)
Recombination factor
Deposited charge

Energy

- For charged current (CC) interactions, E_ν is the sum of the visible components (E_ℓ, E_h) minus the binding energy of the nucleus E_{nuc}

$$E_\nu = E_\ell + E_h - E_{nuc}$$



$$\Delta x = \int_0^{E_0} \frac{1}{dE/dx} dE$$



$$\theta_0 = \frac{S_2}{p\beta c} z \sqrt{\frac{x}{X_0}} \left[1 + \epsilon \log \left(\frac{x}{X_0} \right) \right]$$

$$E_{\text{shower}} = \underbrace{W_i}_{[MeV]} \cdot \underbrace{C_{\text{calo}}}_{[MeV/e]} \cdot \underbrace{C_{\text{adj}}}_{[e^-/ADC]} \cdot \frac{1}{R} \cdot \sum_Q e^{\frac{t_{\text{drift}}}{\tau}} \underbrace{Q}_{[ADC]}$$

Calorimetric constant (TPC gain)
Recombination factor
Deposited charge

Limitations of LArTPC / SBND


- Slow drift time, about 1 m/ms due to dense medium
 - Cosmic ray in the readout window (SBND on surface) 5-10 / event
 - Use optical information to identify neutrino interaction
- High A target - 40 nucleons
 - Limitations due to unobservable nuclear effects obscure E_ν measurement
 - High particle multiplicity at vertex
 - Secondary hadronic interactions
- π^\pm/μ have similar energy loss profile
- Missing neutral particles




Hierarchical Neutrino Event Reconstruction

Relevant goals of pattern recognition in a detailed LArTPC image...

1. Separate topologically distinguishable **types of activity**
2. Identify **important points** (vertex, start points, end points)
3. Cluster individual **particles** (tracks and full showers)
4. Cluster **interactions**, identify **particle properties** in context



SPINE (Scalable Particle Imaging with Neural Embeddings) [4]




[4] F. Drielsma, K. Terao, L. Dominé, D.H. Koh <https://arxiv.org/abs/2102.01033>

SPINE




SLAC: 2019
 ICARUS: 2020
 SBND: 2022
 2x2/ND: 2023
 DUNE FD: 2025



2D -> 3D Projection

Consume 2D hits in each of 3 projections to make space points

- Finds pairs of hits compatible within a time threshold
- Forms a **space point** from 3 wires where 2 hits are compatible in time to form candidate space point



T. Usher




-  Position considered By cluster3d from combinatorial analysis
-  2D Hit in wire planes from track


- False hits create ghost points, which are de-ghosted using a UResNet CNN

Deghosting

- Label each point in simulation sample as ghost/non-ghost
- Learn ghost/non-ghost from reconstructed spacepoint information



Raw Charge




Ghost vs Non-ghost label

UResNet Sparse CNN

- Point-wise feature extraction uses Sparse Convolutional Neural Network (CNN)
- UResNet architecture used as backbone feature extractor

Input


Output



L. Dominé et al.




Deghosting



Raw Charge

Deghosted Charge

Deghosting





SBND MPVMPPR Test Sample


Deghosted Charge

Semantic Segmentation

- **Track** - Ionizing, rectilinear
- **Shower** - EM shower
- **Michel** - electron from muon decay
- **Delta** - electron from track interaction
- **LE** - amorphous blips




Neutrino + Cosmics





Deghosted Charge

Semantic Segmentation



- **Track** - Ionizing, rectilinear
- **Shower** - EM shower
- **Michel** - electron from muon decay
- **Delta** - electron from track interaction
- **LE** - amorphous blips




Neutrino + Cosmics

SBND MPVMPPR Test Sample

PPN




- Point proposal network (PPN) learns attention mask for points of interest
- L2 and CE losses used to learn point location to sub-voxel resolution



L. Dominé et al.




PPN






Missing point

Extra point



Track with start/end point




Neutrino + Cosmics

SBND MPVMPR Test Sample

Fragment Clustering


- Construct intermediate representation of particles called **fragments**
 - Reduces complexity of downstream GNNs
- GraphSPICE is “smart” version of DBSCAN
 - Avoids clustering particles together at vertices



D. Koh et al.






Fragment Clustering



Over-cluster->

Under-cluster->

ARI = Adjusted Rand Index




Neutrino + Cosmics

SBND MPVMMPR Test Sample

Particle Clustering


- Aggregate fragments into **particles** using a Graph Neural Network (GNN)
- Edge representation is the correlation between fragments



Particle Clustering





ARI = Adjusted Rand Index



SBND Simulation
Work in progress

Neutrino + Cosmics




SBND MPVMPT Test Sample

Interaction Clustering




- Aggregate particles into **interactions** using a Graph Neural Network (GNN)
- Edge representation is the correlation between particles

Particles
~~Fragments~~



Interaction Clustering

ARI = Adjusted Rand Index






Neutrino + Cosmics

SBND MPVMPPR Test Sample

Particle ID

- Photon
- Electron
- Muon
- Pion
- Proton



Predicted class	Proton	0.002 (257)	0.000 (8)	0.007 (177)	0.064 (2,178)	0.872 (93,043)
	Pion	0.001 (141)	0.001 (39)	0.043 (1,156)	0.843 (28,590)	0.113 (12,076)
	Muon	0.002 (191)	0.001 (25)	0.950 (25,821)	0.084 (2,860)	0.013 (1,352)
	Electron	0.058 (6,740)	0.958 (26,157)	0.001 (25)	0.005 (159)	0.001 (76)
	Photon	0.937 (109,491)	0.039 (1,074)	0.000 (7)	0.004 (144)	0.001 (99)
	True class	Photon	Electron	Muon	Pion	Proton


MPVMPPR SBND Simulation Work in Progress

Neutrino + Cosmics


SBND MPVMPPR Test Sample

Particle ID

- Electron / photon separation is crucial for ν_e appearance oscillation searches
- Pi / mu separation achievable using GNN



BNB Neutrino (ν_μ dominant)




Intrinsic ν_e

Flash Matching

- Goal - Find interactions compatible with PMT flashes during beam
- Do this by matching hypothesis flash with PMT flash

SBND Preliminary



Flash Matching




[3] D. Garcia-Gamez, P. Green, A.M. Szelc [Eur. Phys. J. C 81, 349 \(2021\)](#)

- Compute hypothesis using semi-analytical model [3]
- Flash score calculated using χ^2


$$\chi^2 = \frac{1}{N_{PDS}} \sum_i \frac{(R_i - H_i)^2}{R_i}$$

- Purity = neutrinos selected / total selected
- Efficiency = true neutrinos selected / true neutrinos

Purity = 85%
Efficiency = 87%



SPINE (Scalable Particle Imaging with Neural Embeddings) [4]



[4] F. Drielsma, K. Terao, L. Dominé, D.H. Koh <https://arxiv.org/abs/2102.01033>




Analyses

ν_μ CC Selections




- SBND ν_μ CC selections will be used for flux and cross section constraints

Run 18255 Subrun 1 Event 142421
(Cosmics filtered out)



Energy depositions



PID



ν_μ CC Selections



- SBND ν_μ CC selections will be used for flux and cross section constraints
- Similar MicroBooNE Np selection had 49% efficiency and 95% purity [5]
- ICARUS performance shown, SBND has similar results in progress

Purity = true signal events selected / selected events

Efficiency = true signal events selected / true signal events





[5] MicroBooNE, [First Simultaneous Measurement of Differential Muon-Neutrino Charged-Current Cross Sections on Argon for Final States with and without Protons Using MicroBooNE Data](https://arxiv.org/abs/2402.19281) [arxiv:2402.19281](https://arxiv.org/abs/2402.19281)

π^0 Selections

- NC/CC π^0 selections are crucial for shower energy calibration



L. Kasher



π^0 Selections





- NC/CC π^0 selections are crucial for shower energy calibration
- Similar MicroBooNE NC π^0 had 9% efficiency and 53% purity [6]

[6] MicroBooNE, [Measurement of neutral current single \$n_0\$ production on argon with the MicroBooNE detector](#), [arXiv:2205.07943](#)

Purity = true signal events selected / selected events
 Efficiency = true signal events selected / true signal events



$$m_{\pi^0} = 134.98 \text{ MeV}/c^2$$





Conclusion

- Evidence for $\Delta m^2 \sim 1 \text{eV}^2$ neutrino oscillation is being explored by SBN
- LArTPC detector output can naturally be reconstructed using computer vision (CV)
- SPINE is a hierarchical feature extractor for LArTPC using CV
 - CNNs reconstruct point-wise attributes
 - GNNs cluster space points into super structures
 - Post processors are used to do physics (e.g. energy reconstruction, flash matching)
- **Reconstruction is essential to reducing uncertainties and measuring oscillations**
- Early simulation selections show promising performance (Michel electron selection in backups)
- SBND SPINE can reconstruct data reliably
 - Results with data to be shown at upcoming conferences...

Thanks!





Energy

- Can calculate energy of **stopped/contained tracks** in TPC using range based methods

$$\Delta x = \int_0^{E_0} \frac{1}{dE/dx} dE$$

- Match distance Δx to lookup table for E_0

Low momentum ← → High momentum





Energy

- Can calculate energy of **exiting/uncontained tracks** using multiple coulomb scattering (MCS)

$$\theta_0 = \frac{S_2}{p\beta c} z \sqrt{\frac{x}{X_0}} \left[1 + \epsilon \log \left(\frac{x}{X_0} \right) \right]$$

- Fit deflection θ_0 to momentum p




Energy

- Can calculate energy of **showers** by summing up all energy depositions
- Derive calibration constants from data

$$E_{\text{shower}} = W_i \cdot C_{\text{calo}} \cdot C_{\text{adj}} \cdot \frac{1}{R} \cdot \sum_Q e^{\frac{t_{\text{drift}}}{\tau}} Q$$

Work function for Ar (above W_i)
 Shower adjustment factor (above C_{adj})
 Electron lifetime correction (above $e^{\frac{t_{\text{drift}}}{\tau}}$)
 [MeV] (below E_{shower})
 [MeV / e⁻] (below W_i)
 [e⁻ / ADC] (below C_{calo})
 Calorimetric constant (TPC gain) (below C_{calo})
 Recombination factor (below $\frac{1}{R}$)
 Deposited charge (below Q)
 [ADC] (below Q)



Michel e- Selection


- Michel electrons from decaying electrons from muons
- Useful for low energy calibrations
- Similar MicroBooNE Michel had 2% efficiency and ~90% purity [6]



N. Oza





[6] MicroBooNE, Michel Electron Reconstruction Using Cosmic-Ray Data from the MicroBooNE LArTPC [arxiv:1704.02927](https://arxiv.org/abs/1704.02927)



Training Sample

- Multi-particle vertex multi-particle rain (MPVMPR) sample - 2 generators
 - Rain (MPR) \longleftrightarrow **cosmic activity**
 - Vertex (MPV) \longleftrightarrow **neutrino activity**
- Use this over neutrino + cosmic event generators to...
 - Sample particle abundances and energies from uniform distribution
 - Avoid biases from particular neutrino generator



Neutrino +
Cosmics

MPVMPR

Semantic Segmentation




Backbone (L. Dominé)

UResNet (UNet + ResNet + Sparse Conv.) as the **backbone feature extractor**

UNet


- Downsizing -> expand receptive field
- Skip connections -> preserve resolution



ML-based Reconstruction for LArTPCs, F. Drielsma (SLAC)


ResNet

- Identity bypass + convolution -> learns residual transform
- Speeds up learning, enables deeper networks





Sparse Convolutions


- Only applies convolutions on active pixels
- Saves memory, execution speed (dramatically)



Flux



- Constrain flux uncertainties by measuring interactions with well known cross section [4,5]




Cross Section


- Measure cross sections of various neutrino interaction processes

Differing observable final states



Quasi-elastic scattering (QE)





Resonance production (RES)







Deep Inelastic scattering (DIS)



J.A. Formaggio, G. Zeller, Reviews of Modern Physics, 84 (2012)



PID Confusion vs KE




Nue v02




PID Confusion vs KE

- Low energy particles commonly identified as protons





BNB Cosmics Rockbox v02



← Particle abundance MPVMR Test v02

PID Confusion vs Overlap

- This is where electron \rightarrow mu confusion comes from - misclustered track fragments. Probably classified as track in semantic seg stage
- Showers are usually clustered only up to 85% level



Nue v02


Short Baseline Near Detector (SBND)



- First neutrinos recorded on July 3, 2024 - [press release](#)



September 2022 -	December 2022 -	March 2023 -	June 2023 -	September 2023 -	December 2023 -	March 2024 -	June 2024 -	September 2023 -	December 2024 -
------------------	-----------------	--------------	-------------	------------------	-----------------	--------------	-------------	------------------	-----------------



Flash Matching

